

INDUCED ABORTION & PRETERM BIRTHS

Preterm births result in some infant deaths and some disorders, including autism, cerebral palsy, epilepsy and mental retardation.

While there are a over sixty known reasons for preterm births and perhaps others that are unknown, there are now 116 studies from 33 countries indicating induced abortion increases the risk of a preterm birth in a pregnancy subsequent to an abortion. One induced abortion increases the risk by one-third while two almost double the risk. Over 50% of aborting women have two or more abortions.

This risk is acknowledged by the Mayo Clinic, the Institute of Medicine, and many eminent researchers. Those who promote and prosper from killing preborn human beings claim that abortion is completely safe.

Studies showing increased preterm birth risk due to abortion were done in these countries.

- | | | | |
|------------------|------------|----------------|-----------------|
| 1 Australia | 9 France | 17 Netherlands | 25 Singapore |
| 2 Botswana | 10 Germany | 18 Nigeria | 26 Slovenia |
| 3 Brazil | 11 Greece | 19 Norway | 27 South Korea |
| 4 China | 12 Hungary | 20 Pakistan | 28 Sweden |
| 5 Czech Republic | 13 Israel | 21 Poland | 29 Taiwan |
| 6 Denmark | 14 Italy | 22 Romania | 30 Togo |
| 7 Egypt | 15 Iraq | 23 Russia | 31 UK |
| 8 Finland | 16 Japan | 24 Scotland | 32 USA |
| | | | 33 Welsh people |

Information provided by Brent Rooney MSc; <http://www.justiceforkids.webs.com>

Pray that life will be protected at every stage.

Abortion abuses women and kills their babies. Help us help both.

St. Jude's Pro Life Committee

Contact information: 301 598.1638 naughtonjj@juno.com May 2010